

SHORT COMMUNICATION

Notes on *Lagothrix flavicauda* (Primates: Atelidae): oldest known specimen and the importance of the revisions of museum specimens

José Eduardo Serrano-Villavicencio^{1,2}, Luís Fábio Silveira³

¹Programa de Pós-graduação em Mastozoologia, Museu de Zoologia, Universidade de São Paulo. Avenida Nazaré 481, Ipiranga, 04263-000 São Paulo, SP, Brazil.

²Centro de Investigación Biodiversidad Sostenible (BioS), Lima, Peru.

³Museu de Zoologia, Universidade de São Paulo. Avenida Nazaré 481, Ipiranga, 04263-000 São Paulo, SP, Brazil.
Corresponding author: José Eduardo Serrano-Villavicencio (serranovillavicencio@gmail.com)

<http://zoobank.org/B29AF1E9-F78A-475D-AF1F-3AECEBABA626>

ABSTRACT. The yellow-tailed woolly monkey, *Lagothrix flavicauda* (Humboldt, 1812), is a large atelid endemic to the cloud forests of Peru. The identity of this species was uncertain for at least 150 years, since its original description in 1812 without a voucher specimen. Additionally, the absence of expeditions to the remote Peruvian cloud forests made it impossible to collect material that would help to confirm the true identity of *L. flavicauda* during the 19th and first half of the 20th century. Until now, the specimens of *L. flavicauda* collected by H. Watkins, in 1925, in La Lejía (Amazonas, Peru) were thought to be the oldest ones deposited in any scientific collection. Nevertheless, after reviewing the databases of the several international museums and literature, we found one specimen of *L. flavicauda* deposited at the Muséum National d'histoire Naturelle (Paris, France) collected in 1900 by G.A. Baër, in the most eastern part of San Martín (Peru), where the presence of this species was not confirmed until 2011. Thus, Baër's specimen represents the oldest known specimen of the yellow-tailed woolly monkey and the only one coming from the eastern part of the species' distribution. Finally, we highlight the importance of online scientific databases for easily diagnosable species. However, caution needs to be taken when using them. We also discuss the value of scientific collections as sources of new discoveries.

KEY WORDS. Distribution, history, taxonomy, yellow-tailed woolly monkey.

Brief taxonomic history of *Lagothrix flavicauda*

The yellow-tailed woolly monkey, *Lagothrix flavicauda* (Humboldt, 1812), endemic to the Peruvian cloud forest in the Departments of Amazonas, San Martín, Loreto, and Huánuco (Leo Luna 1989, Shantee 2011), is one of the rarest and most threatened primates in the Neotropics. The yellow-tailed woolly monkey is also poorly-represented in scientific collections. For more than 150 years after the original description of the species, nothing was known beyond what was written in specimen's labels and naturalists' field notes.

The taxonomic history of *L. flavicauda* began with Humboldt (1812) who reported "Le choro de la province de Jaen" during his travel with Aimé Bonpland through South America (1799–1803). Humboldt (1812) described this species as a brownish-black monkey, with arms and thighs a tone more obscure than the back, a yellowish-brown face, a prehensile tail shorter than the body, of an olive black and laterally adorned with two yellow stripes. However, no specimen was collected on that ex-

pedition. Humboldt (1812) originally considered *L. flavicauda* as an unknown species of howler monkey (genus *Alouatta*), naming it as *Simia flavicauda*. This author also reported that this species was traded in the region of Jaén and Maynas in the Departments of Cajamarca and Loreto (both in Peru), respectively. Although not supported by any voucher specimen, Humboldt's (1812) description leaves no doubt about the uniqueness of *L. flavicauda* but, based solely on this description, several authors reproduced the erroneous idea that *L. flavicauda* would represent a howler monkey species. Thus, *L. flavicauda* was listed as a member of *Mycetes* (Kuhl 1820, Wagner 1840, Lesson 1840, von Tschudi 1844, I. Geoffroy Saint-Hilaire 1851, Schlegel 1876), *Stentor* (É. Geoffroy Saint-Hilaire 1812, Cuvier 1837), and *Alouatta* (Forbes 1896, Elliot 1913, Tate 1939), being the two former genera junior synonyms of the latter.

The misunderstanding about the real identity of Humboldt's *Simia flavicauda* increased with the description of *Lagothrix* (*Oreonax*) *hendeei* (Thomas 1927a). This same author

claimed that the morphological differences of the monotypic subgenus *Oreonax* deserved a generic distinction from *Lagothrix*, containing *O. hendeii* (Thomas 1927b,c). Thomas (1927a,b,c) based his description on material collected by R.W. Hendee in Puca Tambo (San Martín, Peru) in January 1926. In Thomas' vision, his *Oreonax hendeii* clearly represented a different species of Humboldt's *Simia flavicauda*. Thomas (1927b) concluded that Humboldt's *S. flavicauda* was probably nothing more than "a local *Lagothrix*, perhaps *L. lagothricha*". This taxonomic confusion led Tate (1939: 214, 217) to consider Humboldt's *Simia flavicauda* a composite taxon containing two different species, [*Lagothrix*] *hendeii* and [*Alouatta*] *flavicauda*. On the other hand, Cabrera (1958: 153, 183) stated that Humboldt's *Simia flavicauda* could be a valid but undeterminable species of *Alouatta* and recognized Thomas' *Lagothrix hendeii* as a valid species. Hill (1962: 250) also considered *L. hendeii* as a valid species making no mention to Humboldt's *Simia flavicauda*. Nevertheless, these two latter authors considered that the characters presented by Thomas (1927b,c) to sustain *Oreonax* as full genus had no more than specific value and retained *hendeii* within *Lagothrix*. Fooden (1963) performed the first comprehensive taxonomic revision of the genus *Lagothrix*, finally resolving the identity of Humboldt's *Simia flavicauda*. After reviewing the available

literature and comparing museum specimens, Fooden (1963) found that Humboldt's *Simia flavicauda* and Thomas' *Oreonax hendeii* represented the same species of woolly monkey.

Fooden's (1963) taxonomic arrangement was largely accepted (Napier 1976, Eisenberg and Redford 1999, Nowak 1999), until Groves (2001) tested the phylogeny of the atelids. This latter author, based on a 50% majority rule consensus tree estimated via a parsimony analysis of only 20 cranial characters and an undetermined number of species, found out that *Lagothrix flavicauda* was more closely related to species of *Ateles* than to *L. lagothricha*. These low-supported results (bootstrap support <70%) led Groves to resurrect Thomas' *Oreonax* for *flavicauda*. Groves' (2001) proposal was rapidly accepted (Groves 2005, DeLuycker and Heymann 2007, Rylands and Mittermeier 2009); nevertheless, the use of *Oreonax* as a formal genus for *flavicauda* was also criticized. Matthews and Rosenberger (2008) and Rosenberger and Matthews (2008) replicated and extended Groves' (2001) study and showed that his results were probably a sampling artifact, which undermined the logic of reviving *Oreonax* as a formal genus for *flavicauda*. However, as noted by Pacheco et al. (2009), these authors did not provide a phylogeny that placed *flavicauda* within *Lagothrix*. More recently, molecular evidence has also refuted Groves' (2001) results. Ruiz-Garcia et



Figures 1–4. *Lagothrix flavicauda* (MNHN-ZM-MO-1901-1602) collected by G.A. Baer in September 1900 in the Hacienda Nuevo Loreto (San Martín – Peru): (1) complete skin; (2) close-up of the tail showing the diagnostic yellow-golden hairs surrounding the tail's digital patch; (3) close-up of the face showing the white circumbuccal patch; (4) original Baer's label of the specimen. Photographs provided by Jacques Cuisin (MNHN).

al. (2014) and Di Fiore et al. (2015) presented the first molecular phylogenetic analyses of *Lagothrix flavicauda*. The results of both studies were similar, supporting a monophyletic *Lagothrix* that included *flavicauda* and the other woolly monkey species, agreeing with Fooden's (1963) taxonomic arrangement.

The oldest known specimen of *Lagothrix flavicauda*

As a part of an ongoing study, we were looking for specimens of *Lagothrix flavicauda* deposited in scientific collections or cited in the literature. We found 17 skins and 8 skulls deposited in the following collections: Museo de Historia Natural, Lima, Peru (11 skins, 3 skulls); American Museum of Natural History, New York, USA (AMNH; 2 skins, 2 skulls); Natural History Museum, London, UK (NHM; 3 skins, 3 skulls); and The Museum of Vertebrate Zoology, Berkeley, USA (MVZ; 1 skin). All these specimens have been referenced in the literature somehow. Additionally, Kuhl (1820: 30) stated that one specimen of *Mycetes flavicaudatus* (a synonym of *Lagothrix flavicauda*) was held at the Muséum National d'histoire Naturelle (MNHN, Paris – France). Nonetheless, Elliot (1913: 277) said that this must have been a mistake since he could not find that specimen in the collection and É. Geoffroy Saint-Hilaire (1812) never mentioned it either. Nevertheless, we found one skin at the MNHN, collected in 1900 by G.A. Baër, which until now had not been mentioned in the literature.

Between 1900 and 1901, G.A. Baër traveled throughout South America collecting several animals, most of which were birds. He also collected a considerable number of mammals, including rodents, marsupials, and primates. While in Peru, he traveled to some inhospitable areas between June and December of 1900. One of the specimens collected by Baër was an adult individual of *Lagothrix flavicauda* (Figs 1–4). This specimen was collected at the Hacienda Nuevo Loreto in the Department of San Martín in September 1900 (Simon 1902, Stephens and Traylor Jr 1983). Simon (1902) described this locality as a warm forest region of the Río Mixiollo (= Río Mishollo) valley, three days East of Tayabamba, and at an altitude of 1200 m.a.s.l. (Fig. 5). All of Baër's material was shipped to Paris and housed in the collection of the MNHN. At the MNHN, this specimen was identified as a *Cebus* sp. under the catalog number MNHN-ZM-MO-1901-1602. The skin deposited at this museum represents, chronologically, the oldest known specimen of *L. flavicauda* deposited in a scientific institution.

After 50 years without any records, Mittermeier et al. (1977) registered *L. flavicauda* in Pedro Ruiz Gallo (Amazonas, Peru) and, without knowing Baër's record, they stated that the first specimens collected of *L. flavicauda* were those held at the AMNH. These two specimens were provided by H. Watkins, who collected them at La Lejía (Amazonas, Peru) in April 1925. In 1926, R.W. Hendee, a collector for the Godman-Thomas Expedition to Peru, collected three more specimens, those used by Thomas (1927a, b, c). Based on the available data at that time, Mittermeier et al. (1977) proposed that the distribution range of *L. flavicauda* would encompass the southern part of Amazonas and the mountainous western part of San Martín.




Figure 5. Map showing the historical confirmed records of the *Lagothrix flavicauda* specimens deposited in scientific collections. MUSM (circle). AMNH (square), NHM (triangle), and MNHN (star). For locality data, see Table 1.

The northern limit of the distribution of *L. flavicauda* was confirmed in several studies after its Mittermeier et al.'s (1977) record (Leo Luna 1980, Butchart et al. 1995). Nonetheless, its Southern limit remained uncertain for several decades. N. Shanee et al. (2007), based on information given by local guides, informed that this species inhabits the eastern portion of San Martín (40 km west of Tocache, approximately). A few years later, S. Shanee (2011) confirmed the presence of *L. flavicauda* in eastern San Martín. Recently, Aquino et al. (2016) recorded five groups of *L. flavicauda* on the eastern side of the Huallaga River, between Oso Mayo and Palizada (Huánuco, Peru) expanding its southern limits. Although Baër's specimen remained unnoticed, its importance is remarkable. It is not only the first record of *Lagothrix flavicauda* in eastern San Martín, more than 100 years before S. Shanee's (2011) survey, but also, as far as we know, it is the only specimen coming from that region in any scientific collection. Furthermore, if this specimen had been discovered before, the area where it was collected could have been taken into account in the early conservation programs developed for *L. flavicauda*. For example, this area could have been considered to be protected in the buffer zone of the Río Abiseo National Park created in 1990.

Table 1. Summary of the historical records of *Lagothrix flavicauda*, arranged in latitudinal order.

Locality #	Locality	Coordinates	Source
1	Alva (between Chachapoyas and Bagua Grande), Amazonas	5°56'S 77°56'W	MUSM 41, 42, 43, 44, 45
2	Comunidad Las Higueras, Campamento El Triunfo, Amazonas	5°37'S 72°17'W	MUSM 23155
3	La Lejía, Amazonas	6°10'S 77°31'W	AMNH 73222, 73223
4	Puca Tambo, Amazonas	6°10'S 77°16'W	BMNH 27.1.1.1, 27.1.1.2
5	Hacienda Nuevo Loreto, San Martín	8°15'S 76°52'W	MNHN-ZM-MO-1901-1602

The importance of reviewing museum specimens

The specimen of *Lagothrix flavicauda* collected by Baër represents a good example of the importance of thoroughly revising specimens in scientific collections. There are notable examples of new species that were described based on museum material long after the type specimens were collected: *Pithecia vanzolinii* Hershkovitz, 1987, *Callicebus vieirai* Gualda-Barros, Nascimento & Amaral, 2012 and *Mico marcai* (Alperin, 1993).

Vanzolini's bald-faced saki, *P. vanzolinii*, was described based on a large set of samples that remained unnoticed in at least three different Brazilian collections: the Museu de Zoologia da Universidade de São Paulo, São Paulo (MZUSP); the Museu Nacional Universidade Federal do Rio de Janeiro, Rio de Janeiro (MNRJ); and the Museu Paraense Emílio Goeldi, Belém (MPEG). The majority of samples of *P. vanzolinii* were collected in 1936 by A.M. Olalla and other two samples were collected in 1956 by F. Novaes and M.M. Moreira, held at the MPEG (Nunes and Serrano-Villavicencio, 2017). De Vivo (1985) was the first to notice that the material of *Pithecia* collected by Olalla and held at the MZUSP possessed a particular phenotype; nevertheless, he did not recognize this material as a different taxon. It was not until Hershkovitz (1987) that this species was formally described, originally as a subspecies of *P. irrorata*.

Another example is Gualda-Barros et al.'s (2012) *C. vieirai*. The holotype and the paratypes of Vieira's titi were collected in 1997 by Marília Kerr and in 1988 by J.L. Silva-Filho, respectively. All these specimens were kept in the collection of the MZUSP. Gualda-Barros et al. (2012), after revising the collection of *Callicebus* of the MZUSP, noticed that the phenotype of those specimens, deposited there for at least 15 years, was different, and described it as a new species.

As a final example, Marca's Marmoset, *M. marcai*, was described using material collected by the Comissão Rondon in 1914 and held at the MNRJ. This material remained unnoticed in various revisions of callithricids (Hershkovitz 1975, 1977, Ávila-Pires 1969, de Vivo 1991); Alperin (1993) noticed that these specimens represented a different taxon describing it as *Callithrix argentata marcai*. Van Roosmalen et al. (2000) described the marmoset *Mico manicorensis* (Roosmalen, Roosmalen, Mittermeier & Rylands, 2000); nevertheless, the authors did not analyze the type material of *M. marcai* and this later species was found to be a synonym of the former (Garbino 2014).

This kind of 'museum discovery' it is not unusual, but the lack of thorough reviews of scientific collections, especially the

old ones, is alarming and may also have important consequences for conservation, as in the case of critically endangered species. The revision of scientific material brings not only new data and even new species in the middle of the present biodiversity crisis, but also has a tremendous potential to discover new or historical localities which must be viewed as opportunities to evaluate the real conservation status of a species over time (Graham et al. 2004). These historical distributional records provide unique opportunities to trace distributional changes in relation to threatening processes and thereby anticipate future impacts (Drost and Fellers 1996, Shaffer et al. 1998, Ponder et al. 2001, Graham et al. 2004).

Scientific collections have proved to be a source of new discoveries (Funk 2018); in this context, the efforts of all the museums worldwide to make their collections available in digital media should be highlighted and celebrated. However, we recommend that specialists, whenever possible, review these collections carefully to avoid identification mistakes and the posterior uncontrolled misuse of the museum's data. Finally, a proper maintenance of these collections must be guaranteed for future generations of scientists to have access to the valuable material and unfinished source of information that is deposited in them.

ACKNOWLEDGEMENTS

We would like to thank especially Jacques Cuisin from the MNHN, Paris, for kindly providing the information and images needed for this work; Cindy Hurtado for editing the images and for the comments; Victor Pacheco for the access to the database of the Museo de Historia Natural (Lima); João Pedro Souza-Alves and Guilherme Garbino for the valuable comments of the manuscript. Fabio O. do Nascimento and the another anonymous reviewer for the valuable comments on the manuscript. Finally, we thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES – Finance Code 001) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq 302291/2015-6) for the grants to JESV and LFS, respectively.

LITERATURE CITED

Alperin R (1993) *Callithrix argentata* (Linnaeus, 1771): considerações taxonômicas e descrição de subespécie nova. Boletim do Museu Paraense Emílio Goeldi Série Zoologia 9(2): 317–328.

- Aquino R, Charpentier E, García G, López L (2016) First Record of *Lagothrix flavicauda* on the eastern side of the Río Huallaga: An expansion of its known geographic distribution. *Primate Conservation* 30: 15–20.
- Ávila-Pires FD (1969) Taxonomia e zoogeografia do gênero *Callithrix* (Erxleben, 1777) (Primates: Callitrichidae). *Revista Brasileira de Biologia* 29: 49–64.
- Butchart SM, Barnes R, Davies CWN, Fernandez M, Seddon N (1995) Observations of two threatened primates in the Peruvian Andes. *Primate Conservation* 16: 15–19.
- Cabrera A (1958) Catálogo de los mamíferos da América del Sur. *Revista del Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”*, Ciencia Zoológica, Buenos Aires 4: 1–307.
- Cuvier G (1837) The animal kingdom, arranged according to its organization, serving as a foundation for the natural history of animals: and an introduction to comparative anatomy. Volume I. G. Henderson, 1834–37. London, 408 pp.
- DeLuycker AM, Heymann EW (2007) Peruvian yellow-tailed woolly monkey *Oreonax flavicauda* (Humboldt, 1812). In: *Primates in peril: the world’s 25 most endangered primates, 2006–2008*, R. A. Mittermeier et al. (eds.), pp.20–21, 32. *Primate Conservation* (22): 1–40.
- De Vivo M (1985) On some monkeys from Rondônia, Brazil (Primates: Callitrichidae, Cebidae). *Papéis Avulsos de Zoologia* 36(11): 103–110.
- De Vivo M (1991) Taxonomia de *Callithrix* Erxleben, 1777 (Callitrichidae, Primates). *Fundação Biodiversitas para Conservação da Diversidade Biológica*, Belo Horizonte, Brazil, 105 pp.
- Di Fiore A, Chaves PB, Cornejo FM, Schmitt CA, Shantee S, Corté-Ortiz L, Fagundes V, Roos C, Pacheco V (2015) The rise and fall of a genus: Complete mtDNA genomes shed light on the phylogenetic position of yellow-tailed woolly monkeys, *Lagothrix flavicauda*, and on the evolutionary history of the family Atelidae (Primates: Platyrrhini). *Molecular Phylogenetics and Evolution* 82: 495–510. <https://doi.org/10.1016/j.ympev.2014.03.028>
- Drost CA, Fellers GM (1996) Collapse of a regional frog fauna in Yosemite area of the California Sierra Nevada USA. *Conservation Biology* 10: 414–425.
- Eisenberg JF, Redford KH (1999) *Mammals of the Neotropics*. Chicago, The University of Chicago Press, vol. 3, 609 pp.
- Elliot DG (1913) [dated 1912]. *A Review of the Primates*. New York, American Museum of Natural History, vol. 1, 596 pp.
- Fooden J (1963) A revision of the Woolly Monkeys (Genus *Lagothrix*). *Journal of Mammalogy* 44: 213–247. <https://doi.org/10.2307/1377454>
- Forbes HO (1896) *A hand-book to the primates*. London, Edward Lloyd, vol. 1, xiv+286 pp.
- Funk VA (2018) Collections based science in the 21st Century. *Journal of Systematics and Evolution* 56: 175–193. <https://doi.org/10.1111/jse.12315>
- Garbino GST (2014) The Taxonomic Status of *Mico marcai* (Alperin 1993) and *Mico manicorensis* (van Roosmalen et al. 2000) (Cebidae, Callitrichinae) from Southwestern Brazilian Amazonia. *International Journal of Primatology* 35(2): 529–546. <https://doi.org/10.1007/s10764-014-9766-4>
- Geoffroy Saint-Hilaire É (1812) *Tableau des quadrumanes ou des animaux composant le premier ordre de la class des mammifères*. *Annales du Muséum d’histoire naturelle* 19: 85–122.
- Geoffroy Saint-Hilaire I (1851) *Catalogue méthodique de la collection des mammifères, de la collection des oiseaux et des collections annexes, Part. 1 (Catalogue des Primates)*. Paris, Gide et Baudry, Libraires-Éditeurs, xv+96 pp.
- Graham CH, Ferrier S, Huettman F, Moritz C, Peterson AT (2004) New developments in museum-based informatics and applications in biodiversity analysis. *Trends in Ecology & Evolution* 19: 497–503. <https://doi.org/10.1016/j.tree.2004.07.006>
- Groves CP (2001) *Primate Taxonomy*. Washington DC, Smithsonian Institution Press, 350 pp.
- Groves CP (2005) Order Primates. In: Wilson DE, Reeder DM (Eds) *Mammal Species of the World: A Taxonomic and Geographic Reference*. Baltimore, John Hopkins University Press, 3rd ed., 111–184.
- Gualda-Barros J, Nascimento FO, Amaral MK (2012) A new species of *Callicebus* Thomas, 1903 (Primates, Pitheciidae) from the states of Mato Grosso and Pará, Brazil. *Papéis Avulsos de Zoologia* 52(23): 261–279. <https://doi.org/10.1590/S0031-10492012002300001>
- Hershkovitz P (1975) Comments on the taxonomy of Brazilian marmosets (*Callithrix*, Callitrichidae). *Folia Primatologica* 24: 137–172.
- Hershkovitz P (1977) *Living New World Monkeys (Platyrrhini): With an Introduction to Primates*. Chicago, The University of Chicago Press, vol.1, 1132 pp.
- Hershkovitz P (1987) The taxonomy of South American sakis, genus *Pithecia* (Cebidae, Platyrrhini): a preliminary report and critical review with the description of a new species and a new subspecies. *American Journal Primatology* 12(4): 387–468.
- Hill W (1962) *Primates. Comparative Anatomy and Taxonomy*. IV. Cebidae Part A. Edinburgh, Edinburgh University Press, 537 pp.
- Humboldt A (1812) *Tableau synoptique des singes de l’Amérique*. In: Humboldt A. and Bonpland, A. *Recueil d’observations de zoologie et d’anatomie comparée. Faites Dan l’océan Atlantique, dans l’interieur du nouveocontinent at dans la mer de Sud pedant les annés 1799–1803*. Premier volume Deuxieme partie. Observations de zoologie et d’anatomie comparee. Paris, Schoell and Dofour & Co, 368 pp.
- Kuhl H (1820) *Beiträge zur Zoologie und vergleichenden Anatomie*. Frankfurt am Main, Hermannschen Buchhandlung, 212 pp.
- Leo Luna M (1980) First field study of the yellow-tailed woolly monkey. *Oryx* 15: 386–389.

- Leo Luna M (1989) Biología y conservación del mono choro de cola amarilla (*Lagothrix flavicauda*), especie en peligro de extinción. In: Saavedra CJ, Mittermeier RA, Santos IB (Eds) La Primatología en Latinoamérica. Washington DC, World Wildlife Fund, 23–30.
- Lesson RP (1840) Species des mammifères: Bimanés et quadrumanes. París, J.B. Balliere, xiv+292 pp.
- Matthews LJ, Rosenberger AL (2008) Taxon combinations, parsimony analysis (PAUP*), and the taxonomy of the Yellow-Tailed Woolly Monkey, *Lagothrix flavicauda*. American Journal of Physical Anthropology 137(3): 245–55. <https://doi.org/10.1002/ajpa.20859>
- Mittermeier RA, de Macedo-Ruiz H, Luscombe BA, Cassidy J (1977) Rediscovery and conservation of the Peruvian yellow-tailed woolly monkey (*Lagothrix flavicauda*). In: Prince Rainier III, Bourne GH (Eds) Primate Conservation. New York, Academic Press, 95–116.
- Napier PH (1976) Catalogue of Primates in the British Museum (Natural History). Part I: Families Callitrichidae and Cebidae. London, British Museum (Natural History), 121 pp.
- Nowak R (1999) Walker's Mammals of the World, Baltimore, The Johns Hopkins University Press, vol. 1, 642 pp.
- Nunes A, Serrano-Villavicencio J (2017) Rediscovery of Vanzolini's Bald-Faced Saki, *Pithecia vanzolinii* Hershkovitz, 1987 (Primates, Pitheciidae): first record since 1956. Check List 13(1): 2048. <https://doi.org/10.15560/13.1.2048>
- Pacheco V, Cadenillas R, Salas E, Tello C, Zeballos H (2009) Diversidad y endemismo de los mamíferos del Perú. Revista Peruana de Biología 16(1): 5–32. <https://doi.org/10.15381/rpb.v16i1.111>
- Ponder WF, Carter GA, Flemons P, Chapman RR (2001) Evaluation of museum collection data for the use in biodiversity assessment. Conservation Biology 15: 648–657. <https://doi.org/10.1046/j.1523-1739.2001.015003648.x>
- Rosenberger A, Matthews L (2008) *Oreonax*- not a genus. Neotropical Primates 15(1): 8–12. <https://doi.org/10.1896/044.015.0102>
- Ruiz-García M, Pinedo-Castro M, Shostell JM (2014) How many genera and species of woolly monkeys (Atelidae, Platyrrhine, Primates) are there? The first molecular analysis of *Lagothrix flavicauda*, an endemic Peruvian primate species. Molecular Phylogenetics and Evolution 79: 179–198. <https://doi.org/10.1016/j.ympev.2014.05.034>
- Rylands AB, Mittermeier RA (2009) The Diversity of the New World Primates (Platyrrhini). In: Garber PA, Estrada A, Bicca-Marques JC, Heymann EW, Strier KB (Eds) South American Primates: Comparative Perspectives in the Study of Behavior, Ecology, and Conservation. New York, Springer, 23–54. https://doi.org/10.1007/978-0-387-78705-3_2
- Schlegel H (1876) Les singes: Simiae. Leiden, Muséum d'Histoire Naturelle du Pays-Bas, vol. 7, monograph 40, 356 pp.
- Shaffer HB, Fisher R, Davidson C (1998) The role of natural history collections in documenting species declines. Trends in Ecology & Evolution 13: 27–30. [https://doi.org/10.1016/S0169-5347\(97\)01177-4](https://doi.org/10.1016/S0169-5347(97)01177-4)
- Shanee N, Shanée S, Maldonado AM (2007) Conservation assessment and planning for the yellow tailed woolly monkey (*Oreonax flavicauda*) in Peru. Wildlife Biology in Practice 3: 73–82.
- Shanee S (2011) Distribution survey and threat assessment of the yellow-tailed woolly monkey (*Oreonax flavicauda*; Humboldt 1812), Northeastern Peru. International Journal of Primatology 32: 691–707. <https://doi.org/10.1007/s10764-011-9495-x>
- Simon E (1902) Étude sur les Trochilidés observés au Pérou par G. A. Baer (1900–1901). Novitates Zoologicae. Band 9, S. 177–183.
- Stephens L, Traylor Jr MA (1983) Ornithological gazetteer of Peru. Cambridge, Museum of Comparative Zoology, Harvard University, 271 pp.
- Tate GHH (1939) The mammals of the Guiana region. Bulletin of the American Museum of Natural History 76: 151–229.
- Thomas O (1927a) A remarkable new monkey from Peru. The Annals and Magazine of Natural History 9(19): 156–157.
- Thomas O (1927b) The Godman-Thomas Expedition to Peru – V. On mammals collected by Mr. R. W. Hendee in the Province of San Martín, N. Peru, mostly at Yurac Yacu. The Annals and Magazine of Natural History 9(19): 361–375.
- Thomas O (1927c) The Godman-Thomas Expedition to Peru – VI. On mammals from the upper Huallaga and neighbouring highlands. The Annals and Magazine of Natural History 9(20): 594–608.
- Van Roosmalen M, van Roosmalen T, Mittermeier RA, Rylands AB (2000) Two new species of marmoset, genus *Callithrix* Erxleben, 1777 (Callitrichidae, Primates) from the Tapajós/Madeira interfluvium, south central Amazonia, Brazil. Neotropical Primates 8: 2–18.
- Von Tschudi JJ (1844) Untersuchungen über die Fauna Peruana. Part 1 (Therologie). Scheitlin and Zollikofer, St. Gallen, xxx+262 pp.
- Wagner JA (1840) Schreber's Die Säugthiere. Suppl. Vol. 1. Expedition des Schreber'schen Säugthier- und des Esper'schen Schmetterlingswerkes, etc. Erlangen, xiv+551 pp.

 Submitted: September 24, 2018

Accepted: December 11, 2018

Available online: July 5, 2019

Editorial responsibility: Diego Astúa de Moraes

 Author Contributions: JESV and LFS contributed equally to this article.

Competing Interests: The authors have declared that no competing interests exist.

 © 2019 Sociedade Brasileira de Zoologia. Published by Pensoft Publishers at <https://zoologia.pensoft.net>